Queensland Government

Ureterolithotomy (Open)

Facility:

	(Allix identificat	ion label fiele)		
URN:				
Family name:				
Given name(s):				
Address:				
Date of birth:		Sex: M	□F	

(Affix identification label here)

A. Interpreter / cultural needs	
An Interpreter Service is required?	Yes No
If Yes, is a qualified Interpreter present?	Yes No
A Cultural Support Person is required?	Yes No
If Ves is a Cultural Support Person present?	Ves No

B. Condition and treatment

The doctor has explained that you have the following condition: (Doctor to document in patient's own words)

This condition requires the following procedure. (Doctor to document - include site and/or side where relevant to the procedure)

The following will be performed: The removal of a stone from the

The ureter is the tube that drains urine from the kidney to the

C. Risks of a ureterolithotomy (open)

There are risks and complications with this procedure. They include but are not limited to the following.

General risks:

- Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Aspirin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

Specific risks:

- A leak of urine can develop from the cut in the ureter. This may leak on to the skin surface and may continue for some time.
- A narrowing of the tube may develop at the site of where the stone was removed. This may cause damage to the kidney.
- The stone or part of it may not be found and may move down to the bladder or back to the kidney. The stone or fragment may not be able to be removed.
- Deep bleeding in the abdominal cavity. This may need fluid replacement or further surgery.

- Infections such as pus collections in the abdominal cavity. This may need surgical drainage and antibiotic therapy.
- Especially in a male, there may be difficulty in passing urine. A tube (catheter) may need to be put into the bladder to drain the urine. This is usually temporary until the bladder recovers.
- The bowel may be paralysed or blocked after surgery. This may cause a build-up of fluid in the bowel with bloating and vomiting. This may need further treatment.
- A weakness in the wound resulting in the wound bursting in the short term. A hernia may develop in the long term requiring further surgery.
- In some people, healing of the wound may be abnormal and the wound can be thickened and red and the scar may be painful.
- Adhesions (bands of scar tissue) may form and cause a bowel blockage. This can be a short term or long term complication and may need further surgery.

D.	Significant	risks	and	procedure	options
----	--------------------	-------	-----	-----------	---------

(Doctor to document in space provided. Continue in Medical Record if necessary.)
E. Risks of not having this procedure
(Doctor to document in space provided. Continue in Medical Record if necessary.)

	Λ		41	4: _
_	Δn	aesi	me	TIC:

This procedure may require an anaesthetic. (Doctor to document type of anaesthetic discussed)

G. Patient consent

I acknowledge that the doctor has explained;

- my medical condition and the proposed procedure, including additional treatment if the doctor finds something unexpected. I understand the risks, including the risks that are specific to me.
- the anaesthetic required for this procedure. I understand the risks, including the risks that are specific to me.

DURAL CONSENT FC

	Queensland Government
THE T	Government

Facility:

Ureterolithotomy (Open)

	(Affix identification la	bel here)		
URN:				
Family name:				
Given name(s):				
Address:				
Date of birth:		Sex: M	□F	

 other relevant procedure/treatment options and their associated risks.

- my prognosis and the risks of not having the procedure.
- that no guarantee has been made that the procedure will improve my condition even though it has been carried out with due professional care.
- the procedure may include a blood transfusion.
- tissues and blood may be removed and could be used for diagnosis or management of my condition, stored and disposed of sensitively by the hospital.
- if immediate life-threatening events happen during the procedure, they will be treated based on my discussions with the doctor or my Acute Resuscitation Plan.
- a doctor other than the consultant/specialist may conduct/assist with the clinically appropriate procedure/treatment/investigation/examination. I understand this could be a doctor undergoing further training. I understand that all surgical trainees are supervised according to relevant professional guidelines.

I was able to ask questions and raise concerns with the doctor about my condition, the proposed procedure and its risks, and my treatment options. My questions and concerns have been discussed and answered to my satisfaction.

I understand I have the right to change my mind at any time, including after I have signed this form but, preferably following a discussion with my doctor.

I understand that image/s or video footage may be recorded as part of and during my procedure and that these image/s or video/s will assist the doctor to provide appropriate treatment.

Student examination/procedure for educational purposes

For the purpose of undertaking professional training, a student/s may observe the medical examination/s or procedure/s and may also, subject to patient consent, perform an examination/s or assist in performing the procedure/s on a patient while the patient is under anaesthetic. This is for education purposes only. A student/s who undertakes an examination/s or assists in performing the procedure/s will be under the supervision of the treating doctor, in accordance with the relevant professional guidelines.

For the purposes of education I consent to a student/s undergoing training to:

	• •	
•	observe examination/s or procedure/s	☐ Yes ☐ No
•	assist and/or perform examination/s	Yes No
	or procedure/s	

Student - this may include medical, nursing, midwifery, allied health or ambulance students.

I have been given the following Patient Information

Officeus.	
	About Your Anaesthetic
	Ureterolithotomy (Open)

On the basis of the above statements,

I request to have the procedure

Name of Patient:
Signature:
Data:

Patients	who lack capacity to provide consent				
	t must be obtained from a substitute decision in the order below.				
Does the (AHD)?	patient have an Advance Health Directive				
☐ Yes ▶	Location of the original or certified copy of the AHD:				
□ No ▶	Name of Substitute Decision Maker/s:				
	Signature:				
	Relationship to patient:				
	Date: PH No:				
	Source of decision making authority (tick one):				
	Tribunal-appointed Guardian				
	Attorney/s for health matters under Enduring Power of Attorney or AHD				
	Statutory Health Attorney				
	If none of these, the Adult Guardian has provided				

H. Doctor / delegate statement

I have explained to the patient all the above points under the Patient Consent section (G) and I am of the opinion that the patient/substitute decision-maker has understood the information.

are memeration.
Name of
Doctor/delegate:
-
Designation:
Signatura
Signature:
Date:
Duto!

I. Interpreter's statement

I have given a sight translation in

(state the patient's language here) of the consent form and assisted in the provision of any verbal and written information given to the patient/parent or guardian/substitute decision-maker by the doctor.

guardian/substitute decision-maker by the doctor.
Name of
Interpreter:
Signature:
Date:



Consent Information - Patient Copy Ureterolithotomy (Open)

1. What do I need to know about this procedure?

A Ureterolithotomy is an open surgical procedure that will remove a stone from the ureter.

The ureter is the tube that drains urine from the kidney to the bladder.

2. My anaesthetic:

This procedure will require an anaesthetic.

See About Your Anaesthetic information sheet for information about the anaesthetic and the risks involved. If you have any concerns, discuss these with your doctor.

If you have not been given an information sheet, please ask for one.

3. What are the risks of this specific procedure?

There are risks and complications with this procedure. They include but are not limited to the following.

General risks:

- · Infection can occur, requiring antibiotics and further treatment.
- Bleeding could occur and may require a return to the operating room. Bleeding is more common if you have been taking blood thinning drugs such as Warfarin, Aspirin, Clopidogrel (Plavix or Iscover) or Dipyridamole (Persantin or Asasantin).
- Small areas of the lung can collapse, increasing the risk of chest infection. This may need antibiotics and physiotherapy.
- Increased risk in obese people of wound infection, chest infection, heart and lung complications, and thrombosis.
- Heart attack or stroke could occur due to the strain on the heart.
- Blood clot in the leg (DVT) causing pain and swelling. In rare cases part of the clot may break off and go to the lungs.
- Death as a result of this procedure is possible.

Specific risks:

- A leak of urine can develop from the cut in the ureter. This may leak on to the skin surface and may continue for some time.
- A narrowing of the tube may develop at the site of where the stone was removed. This may cause damage to the kidney.
- The stone or part of it may not be found and may move down to the bladder or back to the kidney. The stone or fragment may not be able to be removed.
- Deep bleeding in the abdominal cavity. This may need fluid replacement or further surgery.
- Infections such as pus collections in the abdominal cavity. This may need surgical drainage and antibiotic therapy.

- Especially in a male, there may be difficulty in passing urine. A tube (catheter) may need to be put into the bladder to drain the urine. This is usually temporary until the bladder recovers.
- The bowel may be paralysed or blocked after surgery. This may cause a build-up of fluid in the bowel with bloating and vomiting. This may need further treatment.
- A weakness in the wound resulting in the wound bursting in the short term. A hernia may develop in the long term requiring further surgery.
- In some people, healing of the wound may be abnormal and the wound can be thickened and red and the scar may be painful.
- Adhesions (bands of scar tissue) may form and cause a bowel blockage. This can be a short term or long term complication and may need further surgery.

4. Who will be performing the procedure?

A doctor other than the consultant/specialist may conduct/assist with the clinically appropriate procedure/treatment/investigation/examination.

I understand this could be a doctor undergoing further training, and that all trainees are supervised according to relevant professional guidelines.

If you have any concerns about which doctor/clinician will be performing the procedure, please discuss with the doctor/clinician.

For the purpose of undertaking professional training in this teaching hospital, a student/s may observe the medical examination/s or procedure/s.

Subject to your consent, a student/s may perform an examination/s or assist in performing the procedure/s while you are under anaesthetic. This is for education purposes only. A student/s who undertakes an examination/s or assists in performing the procedure/s will be under the supervision of the treating doctor, in accordance with relevant professional guidelines.

If you choose not to consent, it will not adversely affect your access, outcome or rights to medical treatment in any way. You are under no obligation to consent to an examination/s or a procedure/s being undertaken by a student/s for education purposes.

Notes to talk to my doctor about: